arXiv:1407.3467v1 [math.CO] 13 Jul 2014

PROOF OF A CONJECTURE OF MÉSZÁROS AND MORALES ON THE VOLUME OF A FLOW POLYTOPE

JANG SOO KIM

Abstract. We prove a conjecture of Mészáros and Morales on the volume of a flow polytope. Independently from our work, Zeilberger sketched a proof of their conjecture. In fact, our proof is the same as Zeilberger's proof. The purpose of this note is to give a more detailed proof of the conjecture.

1. Introduction

We prove a conjecture of Mészáros and Morales [2] on the volume of a flow polytope, which is a type D analog of the Chan-Robbins-Yuen polytope [1]. Independently from our work, Zeilberger also proved the conjecture and sketched his proof in [5]. In fact, our proof is the same as Zeilberger's proof. The purpose of this note is to give a more detailed proof of the conjecture.

For a function $f(z)$ with a Laurent series expansion at z, we denote by $\mathrm{CT}_z f(z)$ the constant term of the Laurent expansion of $f(z)$ at 0. In other words, if $f(z) = \sum_{n=-\infty}^{\infty} a_n z^n$, then $\mathrm{CT}_z f(z) = a_0$.

The conjecture of Mészáros and Morales can be stated as the following constant term identity.

Conjecture 1. [2, Conjecture 7.6] For an integer $n \geq 2$, we have

$$\mathrm{CT}_{x_n} \mathrm{CT}_{x_{n-1}} \cdots \mathrm{CT}_{x_1} \prod_{j=1}^{n} x_j^{-1}(1-x_j)^{-2} \prod_{1\leq j<k\leq n} (x_k-x_j)^{-1}(1-x_k-x_j)^{-1} = 2^{n^2} \prod_{k=1}^{n} \mathrm{Cat}(k),$$

where $\mathrm{Cat}(k) = \frac{1}{k+1}\binom{2k}{k}$.

Conjecture 1 is a type D_n analog of the constant term identity

$$\mathrm{CT}_{x_n} \mathrm{CT}_{x_{n-1}} \cdots \mathrm{CT}_{x_1} \prod_{j=1}^{n} (1-x_j)^{-2} \prod_{1\leq j<k\leq n} (x_k-x_j)^{-1} = \prod_{k=1}^{n} \mathrm{Cat}(k), \tag{1}$$

which was conjectured by Chan, Robbins, and Yuen [1] and proved by Zeilberger [4].

In Section 2 we recall Zeilberger's reformulation of Morris' constant term identity. In Section 3 we prove Conjecture 1.

2. Zeilberger's reformulation of Morris' constant term identity

Zeilberger [4] rewrote Morris' identity [3] as follows: For nonnegative integers a and b and a positive half integer c, we have

$$\mathrm{CT}_{x_n} \mathrm{CT}_{x_{n-1}} \cdots \mathrm{CT}_{x_1} \prod_{i=1}^{n} (1-x_i)^{-a} x_i^{-b} \prod_{1\leq i<j\leq n} (x_j-x_i)^{-2c} = \frac{1}{n!} \prod_{j=0}^{n-1} \frac{\Gamma(a+b+(n-1+j)c)\Gamma(c)}{\Gamma(a+jc)\Gamma(c+jc)\Gamma(b+jc+1)}. \tag{2}$$

Zeilberger [4] showed that, when $a = 2, b = 0, c = 1/2$, the right hand side of (2) is

$$\frac{1}{n!} \prod_{j=0}^{n-1} \frac{\Gamma(\frac{n+3+j}{2})\Gamma(\frac{1}{2})}{\Gamma(\frac{4+j}{2})\Gamma(\frac{1+j}{2})\Gamma(\frac{2+j}{2})} = \prod_{k=1}^{n} \mathrm{Cat}(k), \tag{3}$$

which together with (2) implies (1).

Date: July 15, 2014.

By Cauchy's integral formula, if $f(z)$ has a Laurent series expansion at 0, we have

$$\mathrm{CT}_z\, f(z) = \frac{1}{2\pi i}\oint_C \frac{f(z)}{z}dz, \tag{4}$$

where C is the circle $\{z : |z| = \epsilon\}$ oriented counterclockwise for a real number $\epsilon > 0$ such that $f(z)$ is holomorphic inside C except 0. Thus (2) can be rewritten as

$$\begin{aligned}\frac{1}{(2\pi i)^n}\oint_{C_n}\cdots\oint_{C_1}\prod_{j=1}^{n}(1-x_j)^{-a}x_j^{-b-1}\prod_{1\le j<k\le n}(x_k-x_j)^{-2c}dx_1\cdots dx_n \\ = \frac{1}{n!}\prod_{j=0}^{n-1}\frac{\Gamma(a+b+(n-1+j)c)\Gamma(c)}{\Gamma(a+jc)\Gamma(c+jc)\Gamma(b+jc+1)},\end{aligned} \tag{5}$$

where C_j is the circle $\{z : |z| = j\epsilon\}$ oriented counterclockwise for a number $0 < \epsilon < \frac{1}{n}$.

3. Proof of the conjecture of Mészáros and Morales

We will prove the following theorem.

Theorem 1. *For a nonnegative integer a and a positive half integer c, we have*

$$\begin{aligned}\mathrm{CT}_{x_n}\mathrm{CT}_{x_{n-1}}\cdots\mathrm{CT}_{x_1}\prod_{j=1}^{n}x_j^{-a+1}(1-x_j)^{-a}\prod_{1\le j<k\le n}(x_j-x_k)^{-2c}(1-x_j-x_k)^{-2c} \\ = 2^{2an+4c\binom{n}{2}-2n}\frac{1}{n!}\prod_{j=0}^{n-1}\frac{\Gamma(a-\frac{1}{2}+(n-1+j)c)\Gamma(c)}{\Gamma(\frac{1}{2}+jc)\Gamma(c+jc)\Gamma(a+jc)}.\end{aligned}$$

We can get Conjecture 1 as the special case $a = 2, c = 1/2$ of Theorem 1 as follows. When $a = 2, c = 1/2$, the right hand side of the formula in Theorem 1 is

$$\frac{2^{n^2+n}}{n!}\prod_{j=0}^{n-1}\frac{\Gamma(\frac{n+2+j}{2})\Gamma(\frac{1}{2})}{\Gamma(\frac{1+j}{2})\Gamma(\frac{1+j}{2})\Gamma(\frac{4+j}{2})}.$$

Since

$$\prod_{j=0}^{n-1}\frac{\Gamma(\frac{n+3+j}{2})\Gamma(\frac{1}{2})}{\Gamma(\frac{4+j}{2})\Gamma(\frac{1+j}{2})\Gamma(\frac{2+j}{2})}\Bigg/\prod_{j=0}^{n-1}\frac{\Gamma(\frac{n+2+j}{2})\Gamma(\frac{1}{2})}{\Gamma(\frac{1+j}{2})\Gamma(\frac{1+j}{2})\Gamma(\frac{4+j}{2})} = \frac{\Gamma(\frac{2n+2}{2})\Gamma(\frac{1}{2})}{\Gamma(\frac{n+2}{2})\Gamma(\frac{n+1}{2})} = 2^n,$$

we obtain Conjecture 1 by (3).

For the rest of this section we prove Theorem 1. From now on we assume that $\epsilon > 0$ is a very small number.

By (4) we have

$$\begin{aligned}&\mathrm{CT}_{x_n}\mathrm{CT}_{x_{n-1}}\cdots\mathrm{CT}_{x_1}\prod_{j=1}^{n}x_j^{-a+1}(1-x_j)^{-a}\prod_{1\le j<k\le n}(x_j-x_k)^{-2c}(1-x_k-x_j)^{-2c} \\ &= \frac{1}{(2\pi i)^n}\oint_{C_n}\cdots\oint_{C_1}\prod_{j=1}^{n}x_j^{-a}(1-x_j)^{-a}\prod_{1\le j<k\le n}(x_k-x_j)^{-2c}(1-x_k-x_j)^{-2c}dx_1\cdots dx_n,\end{aligned}$$

where C_j is the circle $\{z : |z| = j\epsilon\}$ oriented counterclockwise.

Using the change of variables $x_j = \frac{1-z_j}{2}$ or $z_j = 1-2x_j$, the above is equal to

$$\frac{1}{(2\pi i)^n}\oint_{C_n'}\cdots\oint_{C_1'}\prod_{j=1}^{n}\left(\frac{1-z_j}{2}\right)^{-a}\left(\frac{1+z_j}{2}\right)^{-a}$$
$$\times\prod_{1\le j<k\le n}\left(\frac{-z_k+z_j}{2}\right)^{-2c}\left(\frac{z_k+z_j}{2}\right)^{-2c}(-2)^{-n}dz_1\cdots dz_n$$
$$=\frac{(-1)^n 2^{2an+4c\binom{n}{2}-n}}{(2\pi i)^n}\oint_{C_n'}\cdots\oint_{C_1'}\prod_{j=1}^{n}(1-z_j^2)^{-a}\prod_{1\le j<k\le n}(z_j^2-z_k^2)^{-2c}dz_1\cdots dz_n,$$

where C_j' is the circle $\{z : |z-1| = 2j\epsilon\}$ oriented counterclockwise.

Using the change of variables $z_j^2 = y_j$ or $z_j = y_j^{1/2}$, the above is equal to

$$\frac{(-1)^n 2^{2an+4c\binom{n}{2}-2n}}{(2\pi i)^n}\oint_{C_n''}\cdots\oint_{C_1''}\prod_{j=1}^{n}(1-y_j)^{-a}y_j^{-1/2}\prod_{1\le j<k\le n}(y_j-y_k)^{-2c}dy_1\cdots dy_n,$$

where C_j'' is the circle $\{z : |z-1| = 4j\epsilon\}$ oriented counterclockwise. This is because if C_j' is parametrized by $1+2j\epsilon e^{i\theta}$ for $0\le\theta\le 2\pi$, then the image of C_j' under the map $z\mapsto z^2$ can be parametrized by $1+4j\epsilon e^{i\theta}+4j^2\epsilon^2 e^{2i\theta}$ for $0\le\theta\le 2\pi$. Since ϵ is very small we can deform this image to the circle C_j'' without changing the contour integral.

Using the change of variables $t_j = 1-y_j$, the above is equal to

$$\frac{2^{2an+4c\binom{n}{2}-2n}}{(2\pi i)^n}\oint_{C_n'''}\cdots\oint_{C_1'''}\prod_{j=1}^{n}t_j^{-a}(1-t_j)^{-1/2}\prod_{1\le j<k\le n}(t_k-t_j)^{-2c}dt_1\cdots dt_n,$$

where C_j''' is the circle $\{z : |z| = 4j\epsilon\}$ oriented counterclockwise.

Using (5), we finish the proof of Theorem 1.

Acknowledgement

The author would like to thank Dennis Stanton for helpful discussion.

References

[1] C. S. Chan, D. P. Robbins, and D. S. Yuen. On the volume of a certain polytope. *Experiment. Math.*, 9(1):91–99, 2000.

[2] K. Mészáros and A. H. Morales. Flow polytopes of signed graphs and the Kostant partition function. `http://arxiv.org/abs/1208.0140`, to appear in International Mathematics Research Notices.

[3] W. G. Morris. Constant term identities for finite and affine root systems: conjectures and theorems. PhD thesis, University of Wisconsin-Madison, 1982.

[4] D. Zeilberger. Proof of a conjecture of Chan, Robbins, and Yuen. *Electron. Trans. Numer. Anal.*, 9:147–148 (electronic), 1999. Orthogonal polynomials: numerical and symbolic algorithms (Leganés, 1998).

[5] D. Zeilberger. Sketch of a Proof of an Intriguing Conjecture of Karola Mészáros and Alejandro Morales Regarding the Volume of the D_n Analog of the Chan-Robbins-Yuen Polytope (Or: The Morris-Selberg Constant Term Identity Strikes Again!). `http://arxiv.org/abs/1407.2829`

(Jang Soo Kim) Department of Mathematics, Sungkyunkwan University, Suwon 440-746, South Korea
E-mail address: `jangsookim@skku.edu`